

2-19-2013

Dairy Consumption and Hypertension: Arterial Stiffness and Pulse Pressure

Merrill F. Elias

University of Maine - Main, mfelias@maine.edu

Gregory A. Dore

Georgina E. Crichton

Follow this and additional works at: http://digitalcommons.library.umaine.edu/longitudinal_papers



Part of the [Cardiovascular System Commons](#), and the [Nutrition Commons](#)

Repository Citation

Elias, Merrill F.; Dore, Gregory A.; and Crichton, Georgina E., "Dairy Consumption and Hypertension: Arterial Stiffness and Pulse Pressure" (2013). *Maine-Syracuse Longitudinal Papers*. 7.

http://digitalcommons.library.umaine.edu/longitudinal_papers/7

This Letter to the Editor is brought to you for free and open access by DigitalCommons@UMaine. It has been accepted for inclusion in Maine-Syracuse Longitudinal Papers by an authorized administrator of DigitalCommons@UMaine.

Dairy Consumption and Hypertension: Arterial Stiffness and Pulse Pressure Merrill F. Elias, Gregory A. Dore and Georgina E. Crichton

Hypertension. 2013;61:e40; originally published online February 19, 2013;
doi: 10.1161/HYPERTENSIONAHA.111.00733

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2013 American Heart Association, Inc. All rights reserved.
Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://hyper.ahajournals.org/content/61/4/e40>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Hypertension* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Hypertension* is online at:
<http://hyper.ahajournals.org/subscriptions/>

Letter to the Editor

Letters to the Editor will be published, if suitable, as space permits. They should not exceed 500 words (typed double-spaced) plus 5 references in length and may be subject to editing or abridgment.

Dairy Consumption and Hypertension: Arterial Stiffness and Pulse Pressure

To the Editor:

We read with interest the meta-analysis of dairy consumption and hypertension by Soedamah-Muthu et al¹ in the November 2012 issue of *Hypertension*. Using a stringent set of exclusionary criteria, a literature consisting of 1709 unique prospective studies of dairy and milk consumption was reduced to 9 studies yielding an analysis sample size of 52 756 subjects with 15 367 cases of incident hypertension. This well-controlled meta-analysis confirmed hypotheses generated by previous cross-sectional work, that is, total and low-fat dairy were modestly protective with respect to incident hypertension; relative risk=0.97 (95% confidence interval, 0.95–0.99) and 0.96 (95% confidence interval, 0.94–0.98). Pulse wave velocity (PWV), the current gold standard measure of arterial stiffness, was not an outcome measure in this meta-analysis. Not only is PWV now recognized as an important tool in the diagnosis and management of hypertension, it is also an independent predictor of stroke and cardiovascular mortality and morbidity.²

PWV was not excluded from the Soedamah-Muthu et al meta-analysis, nor was it omitted. Our most recent search of the literature (November 27, 2012) indicated no cross-sectional or prospective examination of dairy consumption in relation to this important outcome variable, other than our recent study.³

In a sample of 587 community-based participants of the Maine-Syracuse Study (58% men, mean age 65, SD=12), stroke and dementia excluded, we reported a statistically significant linear cross-sectional relation between self-reported dairy intake, PWV, systolic blood pressure, and pulse pressure, but found no significant association between dairy intake and diastolic blood pressure, total cholesterol, low-density lipoprotein-cholesterol, or high-density lipoprotein-cholesterol. This was true for a basic model adjusting for age, sex, education, and race, as well as an extended model additionally adjusted for height and weight (or waist circumference), heart rate, antihypertensive drug treatment, mean arterial pressure, the lipid values, depressed mood, grains per day, vegetables per day, sweets per day, protein per day, and total food servings per day. Mean PWV values and 95% confidence intervals associated with reported consumption levels of never or seldom, 1 time per week, 2 to 4 times per week, 5 to 6 times per week, and ≥ 7 times per week were as follows: 11.00 (10.3–11.7), 10.8 (10.1–11.5), 10.6 (10.3–11.00), 10.0 (9.6–10.3), and 10.1 (9.8–10.4) for the fully adjusted model and also when, in a final step, self-report of physical activity was added to the model.

Limitations of our study were absence of information on change in blood pressure over time after our self-report dairy consumption survey, and inability to separate high- and low-fat dairy because of

characteristics of our dietary questionnaire. Nevertheless, our study suggests a potentially important next step, that is, longitudinal studies of high- and low-fat dairy and PWV, and inclusion of PWV in future trials. There are 2 important questions: (1) will slowing in the rate of rise over 9 years, now seen for systolic blood pressure in the Alonso et al study,⁴ be paralleled by a slowing in rise in PWV; and (2) does dairy consumption protect against incident-elevated PWV? Recent normative data make it possible to use one or more criteria of incident PWV as an outcome.⁵

Disclosures

None.

Merrill F. Elias

Department of Psychology and
Graduate School of Biomedical Science
University of Maine
Orono, ME

Gregory A. Dore

Department of Psychology
University of Maine
Orono, ME

Georgina E. Crichton

Nutritional Physiology Research Centre
University of South Australia
Adelaide, Australia

1. Soedamah-Muthu SS, Verberne LD, Ding EL, Engberink MF, Geleijnse JM. Dairy consumption and incidence of hypertension: a dose-response meta-analysis of prospective cohort studies. *Hypertension*. 2012;60:1131–1137.
2. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol*. 2010;55:1318–1327.
3. Crichton GE, Elias MF, Dore GA, Abhayaratna WP, Robbins MA. Relations between dairy food intake and arterial stiffness: pulse wave velocity and pulse pressure. *Hypertension*. 2012;59:1044–1051.
4. Alonso A, Steffen LM, Folsom AR. Dairy intake and changes in blood pressure over 9 years: the ARIC study. *Eur J Clin Nutr*. 2009;63:1272–1275.
5. Reference values for Arterial Stiffness' Collaboration. Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'establishing normal and reference values.' *Eur Heart J*. 2010; 31:2338–2350.