

MOLECULAR CLONING, EXPRESSION ANALYSIS AND CHARACTERIZATION OF GRASS CARP IKK-BETA

Haizhou Wang^{1,2}, Chengyu Hu^{1§}

1. College of Life Sciences, Nanchang University, Jiangxi, P.R.China, 330031

2. College of Materials and Chemical Engineering, Pingxiang University, Jiangxi, P.R.China,
330045

ABSTRACT:

IKK-beta is a member of the IKK complex and plays an important role in the innate immune response. IKK-beta can interact with PKR to activate the NF- κ B pathway in response to dsRNA in mammals. In this study, a grass carp (*Ctenopharyngodon idllus*) IKK-beta gene (*CiIKK β*) was obtained by homology cloning and RACE technique. The full length of *CiIKK β* cDNA is 3428 bp, including 185 bp 5'UTR, 906 bp 3'UTR and 2337 bp open reading frame encoding 778 aa. SMART predicts that there is a serine/threonine kinase region at *CiIKK β* protein N-terminal, followed by a UBQ structural domain and a leucine zipper structure (LZ), then a NEMO binding domain (NBD) at C-terminal. Phylogenetic tree showed that *CiIKK β* is highly homologous to zebrafish IKK β (*DrIKK β*). The real-time PCR result showed that the expression of *CiIKK β* was detected in brain, intestine, liver, spleen, kidney, gill and heart tissues. For the purpose of searching for the mechanism of NF- κ B activation mediated by the interaction between *CiIKK β* and grass carp PKR (*CiPKR*), a glutathione S-transferase(GST) pull-down assay was performed. By GST pull down, *CiIKK β* was shown to be physically associated with the N-terminal of *CiPKR*.

KEYWORDS: IKK β , Tissue expression, PKR, Interaction, GST pull down

§Corresponding author. Tel.: +86 79188317270; Fax: +86 79183969530.

E-mail address: hucy2008@163.com