

2001

Evidence of the Kuwaiti Oil Fires in the Dasuopu Glacier Ice Core, Central Himalaya

Shichang Kang

Qin Dahe

Paul Andrew Mayewski
University of Maine - Main, paul.mayewski@maine.edu

Xie Shucheng

Duan Keqin

Follow this and additional works at: https://digitalcommons.library.umaine.edu/ers_facpub

 Part of the [Earth Sciences Commons](#)

Repository Citation

Kang, Shichang; Dahe, Qin; Mayewski, Paul Andrew; Shucheng, Xie; and Keqin, Duan, "Evidence of the Kuwaiti Oil Fires in the Dasuopu Glacier Ice Core, Central Himalaya" (2001). *Earth Science Faculty Scholarship*. 154.
https://digitalcommons.library.umaine.edu/ers_facpub/154

This Letter to the Editor is brought to you for free and open access by DigitalCommons@UMaine. It has been accepted for inclusion in Earth Science Faculty Scholarship by an authorized administrator of DigitalCommons@UMaine. For more information, please contact um.library.technical.services@maine.edu.

The Editor,
Journal of Glaciology

SIR,

Evidence of the Kuwaiti oil fires in the Dasuopu glacier ice core, central Himalaya

During the Persian Gulf War in 1990–91, more than 740 oil wells in Kuwait were sabotaged (Ferek and others, 1992), and more than 600 wells were set afire. The world, and in particular the Middle East, was left with one of the largest man-made environmental disasters, producing a pronounced impact on regional and global climate and environment (Bakan and others, 1991). Oil fire products during the Persian Gulf War migrated over the Northern Hemisphere. The smoke aerosol from long-range transport of Kuwaiti oil fires was observed in Japan (Okada and others, 1992), Pakistan (Limaye and others, 1991), Hawaii (Lowenthal and others, 1992), as well as Wyoming in the U.S.A. (Deshler and Hofmann, 1992).

In summer 1997, a 15 m ice core, covering the period 1988–97, was collected from 7000 m a.s.l. on a relatively flat portion of Dasuopu glacier (28°23' N, 85°44' E) on the north-west margin of Mount Xixiabangma in the central Himalaya. Detailed methods of ice-core sampling and chemical analysis are described by Kang and others (2000), along with a discussion of monsoon and dust signals in the core. Profiles of $\delta^{18}\text{O}$ and major-ion (Ca^{2+} , NO_3^- and SO_4^{2-}) concentrations against

water-equivalent depth are shown in Figure 1. In general, variations of SO_4^{2-} are in agreement with those of Ca^{2+} and NO_3^- in the ice core, mainly reflecting dust input from central Asia (Kang and others, 2000). The highest SO_4^{2-} concentration, 730.7 ppb (marked by an arrow in Figure 1), in the core is recorded during early 1991. The lowest SO_4^{2-} concentration is 6.1 ppb, and mean SO_4^{2-} is 122.9 ppb (std dev. 132.2). Ca^{2+} and NO_3^- concentrations during early 1991 do not have anomalous peaks. Thus, the 1991 SO_4^{2-} is unique and appears not to be related to dust deposition.

In June 1991, the explosive eruption of Mount Pinatubo, Philippines, injected an estimated $(18 \pm 2) \times 10^6$ t of SO_2 directly into the atmosphere (Krueger and others, 1995). The volcanic aerosol mass was dispersed gradually in the global atmosphere, covering the entire Earth by mid-1992 (Hitchman and others, 1995). Pinatubo volcanic signals were identified in South Pole snow, which occurred from early or mid-1992 to mid-1994 (Cole-Dai and Mosley-Thompson, 1999). Though we cannot confirm when volcanic aerosol was transported to the Dasuopu core site, it was probably later than June 1991. However, the highest SO_4^{2-} value in the Dasuopu core is recorded in early 1991 (Fig. 1). Thus, we consider that the 1991 spike of SO_4^{2-} concentration is not related to the Pinatubo eruption.

We assume that the SO_4^{2-} peak in 1991 in the Dasuopu ice core is related to the Kuwaiti oil fire products during the Persian Gulf War. Firstly, the Kuwaiti oil wells were set afire




Fig. 1. Depth profiles of $\delta^{18}\text{O}$ and ionic concentrations with depth (water equivalent) in the Dasuopu ice core. Dating was performed by counting seasonal peaks of $\delta^{18}\text{O}$ and major ions. The coarse solid line is a weighted smoothing (5 points smoothing). Dashed lines indicate annual layers.

in early 1991 (Ferek and others, 1992) and the aerosol smoke from oil fires was transported to the Himalayan region by a southern westerly jet, which only took 1–2 days (Bodhaine and others, 1992). Secondly, measurements of organic matter in the same core indicate that concentrations of some organic compounds from petroleum residues were higher during 1988–92 than in 1997 (Xie and others, 2000). Thirdly, chemical analysis of smoke plumes from the Kuwaiti oil fires shows that SO₂ concentrations are 1–2 orders of magnitude higher than NO_x, and SO₄²⁻ is the dominant species among major ions in smoke plumes (Ferek and others, 1992). These points suggest that pollutants from Kuwaiti oil fires were transported to Dasuopu glacier by the westerly jet and recorded in snow at 7000 m a.s.l. in the Himalaya. Thus, ice-core records from the Himalaya provide a unique opportunity to recover pollutant signals and reconstruct environmental changes in the past.

ACKNOWLEDGEMENTS

This research is supported by the Chinese Academy of Sciences (KZCX2-305, KZ951-A1-402, KZCX2-301/108, KZCX1-10-02), the NKBRFSF Project of China (G1999043400) and the National Nature Science Foundation of China (49871022).

Laboratory of Ice Core and
Cold Regions Environment,
Cold and Arid Regions Environment and
Engineering Research Institute, CAS,
Lanzhou 730000, People's Republic of China

KANG SHICHANG*
QIN DAHE

*Institute for Quaternary
and Climate Studies,
University of Maine,
Orono, Maine 04469-5790, U.S.A.

PAUL A. MAYEWSKI

Laboratory of Ice Core and
Cold Regions Environment,
Cold and Arid Regions Environment and
Engineering Research Institute, CAS,
Lanzhou 730000, People's Republic of China

XIE SHUCHENG
DUAN KEQIN

10 September 2001

REFERENCES

- Bakan, S. A. and 16 others. 1991. Climate response to smoke from the burning oil well in Kuwait. *Nature*, **351**(6325), 367–371.
- Bodhaine, B. A., J. M. Harris, J. A. Ogren and D. J. Hofmann. 1992. Aerosol optical properties at Mauna Loa observatory: long-range transport from Kuwait? *Geophys. Res. Lett.*, **19**(6), 581–584.
- Cole-Dai, J. and E. Mosley-Thompson. 1999. The Pinatubo eruption in South Pole snow and its potential value to ice-core paleovolcanic records. *Ann. Glaciol.*, **29**, 99–105.
- Deshler, T. and D. J. Hofmann. 1992. Measurements of unusual aerosol layers in the upper troposphere over Laramie, Wyoming, in the spring of 1991: evidence for long-range transport from the oil fires in Kuwait. *Geophys. Res. Lett.*, **19**(4), 385–388.
- Ferek, R. J., P. V. Hobbs, J. A. Herring and K. K. Laursen. 1992. Chemical composition of emissions from the Kuwait oil fires. *J. Geophys. Res.*, **97**(D13), 14,483–14,489.
- Hitchman, M. H., M. McKay and C. R. Trepte. 1995. A climatology of stratospheric aerosol. *J. Geophys. Res.*, **99**(D10), 20,698–20,700.
- Kang Shichang, C. P. Wake, Qin Dahe, P. A. Mayewski and Yao Tandong. 2000. Monsoon and dust signals recorded in Dasuopu glacier, Tibetan Plateau. *J. Glaciol.*, **46**(153), 222–226.
- Krueger, A. J. and 6 others. 1995. Volcanic sulfur dioxide measurements from the total ozone mapping spectrometer instruments. *J. Geophys. Res.*, **100**(D7), 14,057–14,076.
- Limaye, S. S., V. E. Suomi, C. Velden and G. Tripoli. 1991. Satellite observations of smoke from oil fires in Kuwait. *Science*, **252**(5012), 1536–1539.
- Lowenthal, D. H., R. D. Borys and J. C. Chow. 1992. Evidence for long-range transport of aerosol from the Kuwait oil fires to Hawaii. *J. Geophys. Res.*, **97**(D13), 14,573–14,580.
- Okada and 6 others. 1992. Kuwait soot over Japan. *Nature*, **355**(6356), 120.
- Xie Shucheng, Yao Tandong, Kang Shichang, Duan Keqin, Xu Baiqin and L. G. Thompson. 2000. Geochemical analysis of a Himalayan snowpit profile: implication for atmospheric pollution and climate. *Organic Chem.*, **31**(1), 15–23.