

The University of Maine

DigitalCommons@UMaine

---

Honors College

---

Spring 5-2017

## The Effects of Mercury Exposure on the Cytochrome C Oxidase 1 Gene of Larval Dragonflies

Megan C. Little  
*University of Maine*

Follow this and additional works at: <https://digitalcommons.library.umaine.edu/honors>



Part of the [Biochemistry Commons](#), and the [Molecular Biology Commons](#)

---

### Recommended Citation

Little, Megan C., "The Effects of Mercury Exposure on the Cytochrome C Oxidase 1 Gene of Larval Dragonflies" (2017). *Honors College*. 436.

<https://digitalcommons.library.umaine.edu/honors/436>

This Honors Thesis is brought to you for free and open access by DigitalCommons@UMaine. It has been accepted for inclusion in Honors College by an authorized administrator of DigitalCommons@UMaine. For more information, please contact [um.library.technical.services@maine.edu](mailto:um.library.technical.services@maine.edu).

THE EFFECTS OF MERCURY EXPOSURE ON THE CYTOCHROME C OXIDASE 1  
GENE OF LARVAL DRAGONFLIES

by

Megan C. Little

A Thesis Submitted in Partial Fulfillment  
of the Requirements for a Degree with Honors  
(Molecular and Cellular Biology/Biochemistry)

The Honors College

University of Maine

May 2017

Advisory Committee:

Sarah Nelson, Associate Research Professor, School of Forest Resources, Advisor  
Keith Hutchison, Professor Emeritus, Molecular and Biomedical Sciences,  
Advisor  
Julie Gosse, Associate Professor of Molecular and Biomedical Sciences  
Michael Kinnison, Professor of Evolutionary Applications, School of Biology and  
Ecology  
Christopher Mares, Director, Intensive English Institute

## ABSTRACT

Mercury is an environmental pollutant; its most toxic form is methylmercury. Once mercury is converted to methylmercury in a body of water it is able to bioaccumulate in organisms and biomagnify up the food chain. Mercury is able to cause DNA damage through the generation of free radicals and binding to sulfhydryl groups of cysteines in zinc finger DNA binding domains, inhibiting DNA repair machinery. In this study the potential mutagenic effects of mercury were investigated on larval dragonflies (Odonata: Anisoptera) collected from national parks across the United States. Since mercury is a known mutagen it was hypothesized that the COI gene from dragonfly larvae collected from sites where they are subjected to higher mercury exposure should have more mutations than the COI gene from dragonfly larvae collected from sites where they are subjected to lower mercury concentrations. The COI gene from each sample was first used to determine species of selected dragonflies through DNA barcoding then was analyzed for mutations in its nucleotide and amino acid sequences. Samples showing mutations in their amino acid sequences were modeled to determine if the mutation caused a change in the protein structure. Mutations were detected that did change protein structure, but at this point it is unknown if this structural change impacts protein function. The mutation rate was ultimately shown not to increase at sites with greater mercury exposure. Instead, the data suggests that genetic variation in dragonflies decreases at higher concentrations of mercury.

## ACKNOWLEDGEMENTS

I would like to thank my thesis advisors, Sarah Nelson and Keith Hutchison, and my committee members, Julie Gosse, Michael Kinnison, and Christopher Mares, for all of their support throughout this process. I would also like to thank Michael Marion and Karen James for providing DNA barcoding expertise, Hamish Greig, Michael Kinnison, and Wesley Wright for providing laboratory support, and Patty Singer and the University of Maine DNA Sequencing Facility for conducting DNA sequencing for these samples. The United States Geological Survey and the National Park Service are core collaborators on the Dragonfly Mercury Project, which provided the dragonfly mercury data used in this thesis. Staff from the NPS, its partners, and over 824 citizen scientists collected samples across the US in 2014. University of Maine faculty research funds to Sarah Nelson and Hamish Greig supported the DNA Barcoding effort.

## TABLE OF CONTENTS

I. Introduction	1
Mercury	1
Pollution	1
Genotoxic Effects	1
Damage Prevention	2
DNA Barcoding	3
Dragonflies	4
II. Materials and Methods	6
Sample Collection and Storage	6
Mercury Analysis	6
DNA Extraction	6
Polymerase Chain Reaction (PCR)	7
DNA Sequencing	7
Sequence Analysis	7
III. Results	8
COI Barcode Validity	8
Nucleotide Sequence Alignments	10
Amino Acid Sequence Alignments	15
Protein Structure Models	16
IV. Discussion and Conclusions	18
Bibliography	23
Appendix	27
Author's Biography	33

## LIST OF FIGURES

Figure 1	5
Figure 2	9
Figure 3	11
Figure 4	12
Figure 5	13
Figure 6	14
Figure 7	15
Figure 8	16
Figure 9	17

## INTRODUCTION

### Mercury

#### Pollution

Mercury is an environmental toxicant with many natural and anthropogenic sources. The largest sources of mercury are oceans and other water bodies, but other natural sources include volcanic eruptions and wildfires.<sup>1, 2, 3</sup> Anthropogenic sources include gold mining and production, fossil fuel combustion, and cement production.<sup>4, 5</sup>

Mercury can be found in many different forms. Mercury emissions into the air take the form of elemental mercury or can be emitted as particulates. Through various oxidation-reduction reactions mercury can form soluble ionic or non-ionic organic or inorganic compounds.<sup>1</sup> The most prevalent form of mercury in the environment is methylmercury.<sup>1, 6</sup> Methylmercury present in aquatic environments is able to bioaccumulate in organisms and biomagnify up the food chain, entering cells by diffusing across cell membranes, due to being soluble in lipids.<sup>7, 8, 9, 10</sup> Once inside the cell, mercury induces a variety of genotoxic effects, such as mutations and chromosomal aberrations. The main ways that mercury causes DNA damage are generation of oxidative stress and interference with DNA repair enzymes.<sup>7</sup>

#### Genotoxic Effects

Oxidative stress is caused by the accumulation of reactive oxygen species (ROS).<sup>7</sup> Mercury allows for the production of ROS because it is easily oxidized by hydrogen peroxide, yielding the hydroxyl free radical.<sup>11, 12, 13</sup> While ROS can directly cause DNA damage, they also cause the generation of the modified nucleotide 8-oxo-deoxyguanosine

triphosphate (8-oxo-dGTP). This nucleotide can be incorporated in genomic or mitochondrial DNA, but the effects are more severe in the mitochondria because its polymerase, polymerase  $\gamma$ , suffers a decrease in fidelity in the presence of 8-oxo-dGTP.<sup>14, 15, 16, 17</sup> ROS can also damage cell membranes, particularly mitochondrial membranes, by inducing autocatalytic lipid peroxidation. In this process, the ROS oxidize a membrane lipid, turning it into an unstable fatty acid radical, prompting its degradation into a lipid hydroperoxide. This process makes the cellular membranes more permeable, and in the mitochondria this makes the electron transport chain less efficient, leading to the production of hydrogen peroxide.<sup>18, 19, 20, 21</sup>

Mercury can also cause an accumulation of DNA mutations due to interference with DNA repair machinery. This is due to the ability of mercury to bind to the sulfhydryl groups of the cysteines present in zinc finger DNA binding domains, which displaces the zinc ion and prevents DNA binding.<sup>22, 23, 24</sup> In a case study involving factory workers occupationally exposed to mercury vapor mercury exposure was shown to negatively impact base excision repair mechanisms more than nucleotide excision repair mechanisms.<sup>22</sup> Mercury can also bind the DNA directly, causing changes in secondary structure which could prohibit DNA machinery binding, or promote/inhibit translation.<sup>7, 25</sup>

### Damage Prevention

Cells produce glutathione (GSH), which acts as an antioxidant and functions to prevent DNA damage from mercury and the resulting ROS it generates. First, GSH contains a sulfhydryl group that allows it to bind to mercury. In this way GSH can sequester mercury compounds before they can be oxidized during the formation of ROS.<sup>26</sup> Second, GSH can neutralize any ROS made using mercury because it is easily

oxidized through the action of glutathione peroxidase.<sup>26, 27, 28</sup> Oxidized GSH can be recycled back to GSH by glutathione reductase in the presence of NADPH.<sup>28</sup>

GSH is produced by  $\gamma$ -glutamylcysteine synthetase and GSH synthase, but there is a feedback mechanism in place where GSH inhibits  $\gamma$ -glutamylcysteine synthetase function.<sup>28, 29, 30, 31</sup> This feedback mechanism is likely in place because if GSH is only partially oxidized, it becomes the glutathione thiol free radical and can increase the production of ROS.<sup>26, 27, 32</sup> GSH has been shown to be depleted in cells exposed to 1000 ppb of mercury, although DNA damage appears at a mercury concentration of only 50 ppb.<sup>26</sup>

### DNA Barcoding

DNA barcoding allows for a species level identification for an organism using a short section of its genome. Identification is based on sequence homology of the section from the sample organism and sections from other organisms whose species was determined by taxonomists and vouchered in an authoritative repository. The Barcode of Life Database (BOLD) is one tool that can be used to make these comparisons (<http://www.boldsystems.org>).

After the initial set up and vouchering required by taxonomists, DNA barcoding is a more reliable way to identify many species for a few reasons. First, phenotypic variation within a population makes taxonomic identification of new or similar species difficult. Secondly, morphological keys used in identification are often specific for only one life stage and/or gender of a species. Lastly, because taxonomic identification requires specialized knowledge of morphological traits, taxonomists can often only identify the species of a small subset of organisms, whereas DNA barcoding is more

accessible.<sup>33</sup> For dragonflies, the adult lifestage is generally well known and described, but different species of larval dragonflies can be difficult to distinguish, in some cases requiring raising of a live larva until it matures.

DNA barcoding can be performed using ribosomal or mitochondrial DNA, however mitochondrial DNA is preferred because it has a higher rate of molecular evolution, allowing for discrimination between closely related species.<sup>34</sup> The mitochondrial cytochrome c oxidase I (COI) gene is currently the favored gene for DNA barcoding of insects like dragonflies because of the other mitochondrial protein-coding genes, COI exhibits the most rapid rate of molecular evolution of its third position nucleotides.<sup>33, 35, 36</sup> COI, along with the cytochrome c oxidase II (COII) gene, forms the final acceptor in the electron transport chain, catalyzing the reduction of oxygen to water.<sup>37</sup>

### Dragonflies

The U.S. Geological Survey (USGS), the University of Maine, and the National Park Service (NPS) are currently involved in the collaborative Dragonfly Mercury Project. For this project, citizen scientists collect larval dragonflies from over 50 national parks, multiple sites per park, and the dragonfly larvae are analyzed for mercury content (Figure 1). In this way dragonflies act as biosentinels for the bioaccumulation of mercury in aquatic food systems.<sup>38</sup>




Figure 1: Parks that participated in the Dragonfly Mercury Project in 2014. Individual water bodies within the parks are separate sampling sites. Most samples in this thesis research were from parks in the eastern portion of the continental U.S.

For this project representative dragonflies from all sample sites were chosen for DNA barcode species identification. The COI sequences of select species were then analyzed for mutations and altered protein structure corresponding to level of mercury exposure. Since mercury is a known mutagen through its involvement in the generation of ROS and interference with DNA repair machinery, the COI gene from dragonfly larvae collected from sites where they are subjected to higher mercury exposure are expected to have more mutations than the COI gene from dragonfly larvae collected from sites where they are subjected to lower mercury concentrations.

## MATERIALS AND METHODS

### Sample Collection and Storage

Dragonflies collected from the field were individually bagged, shipped overnight on dry ice to the University of Maine, and stored at -20°C. 215 samples representing all dragonfly families found at each sampling site were chosen to undergo DNA barcode species identification. The rest were sent to undergo mercury analysis.

### Mercury Analysis

Wet weight of each sample was determined before freeze-drying and determination of dry weight. The University of Maine Sawyer Lab determined total mercury concentrations in dragonfly larvae using a Nippon Instruments Model MA-3000 Mercury Analyzer (Direct Hg by Thermal Combustion, US EPA method 7473). Mercury concentrations of samples found at the same sampling site were averaged to determine geometric mean concentration of mercury exposure in larvae at that site.

### DNA Extraction

Tissue from three of each dragonfly's legs was used for the DNA extraction for each sample. The Qiagen DNeasy Blood and Tissue kit was used for DNA extractions. Leg tissue was combined with Buffer ATL and Proteinase K before being ground and incubated overnight at 55°C. Samples were vortexed, combined with Buffer AL, vortexed again, and incubated at 55°C for 20 minutes. After addition of ethanol, mixture volume was pipetted into a DNeasy Mini Spin Columns and centrifuged at 8000 rpm for 1 minute. Buffer AW1 was added and columns were again centrifuged at 8000 rpm for 1 minute. Buffer AW2 was added and columns were centrifuged at 14,000 rpm for 3

minutes. After placing column into a 1.5 mL microcentrifuge tube, Buffer AE was added, samples incubated at room temperature for 1 minute, and then were centrifuged at 8000 rpm for 1 minute to elute the DNA. Buffer AE was added again, samples were incubated at 55°C for 5 minutes, and then were centrifuged at 8000 rpm for 1 minute to elute the remaining DNA. DNA was quantified using a nanodrop. Purified DNA was stored at -20°C. See part B of Appendix for specific procedure.

#### Polymerase Chain Reaction (PCR)

m1COIintF and jgHCO2198 primers were used for amplification of extracted DNA. Amplified region was 313 bp in length.<sup>34</sup> Master mix contained the following component volumes per reaction: 12.5 µL Sigma RedTaq ReadyMix, 10.5 µL nuclease-free water, 0.5 µL m1COIintF, and 0.5 µL jgHCO2198. 24 µL of master-mix and 1 µL of DNA were used for each reaction. See part C of Appendix for thermal cycler conditions. PCR products were run on 1.5% agarose gels and photographed using a UV transilluminator before being stored at -20°C.

#### DNA Sequencing

PCR products were cleaned using Affymetrix ExoSAP-IT, diluted to a concentration of 10 ng/ µL, and delivered to the University of Maine's DNA sequencing facility.

#### Sequence Analysis

Obtained nucleotide sequences were edited used 4Peaks software. Species was determined using the Barcode of Life Database (BOLD). Amino acid sequence generated using NCBI Blast. Nucleotide and amino acid sequence alignments were performed using Clustal Omega. Protein structure was determined using Phyre2.

## RESULTS

### COI Barcode Validity

Nucleotide and amino acid sequences from five representative species were aligned to test the validity of using COI gene sequences as barcodes for species identification (Figure 2). Nucleotide sequences have a number of variable loci, but due to the redundancy of the genetic code these are reduced in the amino acid sequences.



### Nucleotide Sequence Alignments

Nucleotide sequence alignments were performed on species found at more than one site and with total mercury concentrations ranging from <100 ppb dw (parts per billion, dry weight) to >200 ppb dw to detect possible mutations. Five species were examined (Figures 3 – 7). Each species was shown to have one sample with a mutation.

A

```

65 -----TCTTGCAGGAGCAATTGCACATGCAGGAGCATC
153 -GGTACAGGATGAACTGTTTATCCCCCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
137 -GGTACAGGATGAACTGTTTATCCCCCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
73 TGGTACTGGTTGAACTGTTTATCCCCCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
173 TGGTACAGGTTGAACTGTTTATCCCCCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
208 -----TCTTGCAGGAGCAATTGCACATGCAGGAGCATC
191 -----CCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
178 -----CTTGCAGGAGCAATTGCACATGCAGGAGCATC
84 -----CAGGAGCAATTGCACATGCAGGAGCATC
81 TGGAACAGGATGAACAGTTTATCCCCCTCTTGCAGGAGCAATTGCACATGCAGGAGCATC
*****

65 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTATAGGAGCAAT
153 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
137 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
73 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
173 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
208 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
191 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
178 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
84 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
81 AGTTGATTTAACTATCTTCTCTTTCATCTTGCAGGAGTTTCATCAATTCTAGGAGCAAT
*****

65 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
153 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
137 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
73 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
173 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
208 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
191 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
178 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
84 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
81 TAACTTTATTACTACAATATTAATATAAAAGTCACCAGGAATGAAATTAGACCAAATACC
*****

65 TCTATTGGTATGGGCTGTAATTATTACTGCAGTTTT-----
153 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTAC-----
137 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTAC-----
73 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTATT
173 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTACA
208 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTATT
191 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTATT
178 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTATT
84 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTT-----
81 TCTATTTGTATGGGCTGTAGTTATTACTGCAGTTTTACTTTTATTATCTTTACCAGTATT
*****

```

B

Sample #	Park Code	Site	[Hg] (ppb)	% Sequence Similarity
65	VOYA	Peary Lake	76	98.2
153	ACAD	Long Pond	131	100
137	ACAD	Jordan Pond	145	100
73	CACO	Herring Pond	79	100
173	ACAD	Hodgdon Pond	165	100
208	ACAD	Lakewood	130	100
191	ACAD	Aunt Betty's	234	100
178	ACAD	Eagle Lake	209	100
84	CACO	Great Pond	186	100
81	CACO	Duck Pond	171	100

Figure 3: A. *Gomphus exilis* nucleotide sequence alignments. B. Quantification of part A. Park codes are as follows: VOYA – Voyageurs National Park, CACO – Cape Cod National Seashore, ACAD – Acadia National Park

A

```

59 ACATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGATGGTGTATCATC
40 ACATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
50 ACATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
64 ACATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
88 ACATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
205 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
196 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
149 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
135 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
55 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
54 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
41 -CATGCAGGAGCATCAGTAGATTTAACTATTTTTCTTTTACACTTGGCTGGTGTATCATC
*****

59 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
40 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
50 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
64 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
88 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
205 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
196 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
149 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
135 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
55 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
54 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
41 AATTCTAGGTGCTATTAATTTTATTACTACAACAATTAATATAAAAATCACCAGGAATGAA
*****

59 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
40 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTA-----
50 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
64 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
88 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTAGT
205 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
196 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
149 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTG-----
135 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATT---
55 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
54 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
41 AATAGATCAAATACCTTTATTTGTATGAGCAGTTGTAATTACAGCTGTACTCTTATTACT
*****

```

B

Sample #	Park Code	Site	[Hg] (ppb)	% Sequence Similarity
59	VOYA	Peary Lake	76	99.5
40	PIRO	Beaver Lake	142	100
50	PIRO	Grand Sable Lake	149	100
64	VOYA	Ryan Lake	153	100
88	OZAR	Coppermine Creek	136	100
205	ACAD	Aunt Betty's	234	100
196	ACAD	Jordan Pond	145	100
149	ACAD	Long Pond	131	100
135	ACAD	Jordan Pond	145	100
55	SACN	Earl Landing	135	100
54	SACN	Earl Landing	135	100
41	PIRO	Grand Sable Lake	149	100

Figure 4: A. *Basiaeschna janata* nucleotide sequence alignments. B. Quantification of part A. Park codes are as follows: PIRO – Pictured Rocks National Lakeshore, SACN – Saint Croix National Scenic Riverway, VOYA – Voyageurs National Park, OZAR – Ozark National Scenic Riverway, ACAD – Acadia National Park.

A

```

186 -----CCTCTGGCAGGTACAATTGCCCATGCTGGTTCCTCT
72 -----TCTGGCAGGTACAATTGCCCATGCTGGTTCCTCT
134 GGTACTGGTTGAACAGTTTATCCCCCTCTGGCGGTACAATTGCCCATGCTGGTTCCTCT
85 GGAACAGGATGAACAGTTTATCCCCCTCTGGCAGGTACAATTGCCCATGCTGGTTCCTCT
*****

186 GTAGACCTAACAATTTTCTCTTTACACCTCGCAGGTGTATCATCAATTTTAGGAGCAATT
72 GTAGACCTAACAATTTTCTCTTTACACCTCGCAGGTGTATCATCAATTTTAGGAGCAATT
134 GTAGACCTAACAATTTTCTCTTTACACCTCGCAGGTGTATCATCAATTTTAGGAGCAATT
85 GTAGACCTAACAATTTTCTCTTTACACCTCGCAGGTGTATCATCAATTTTAGGAGCAATT
*****

186 AATTTTATTACCACTGTAATTAACATGAAATCACCAGGTATAAAATTAGACCAAATACCT
72 AATTTTATTACCACTGTAATTAACATGAAATCACCAGGTATAAAATTAGACCAAATACCT
134 AATTTTATTACCACTGTAATTAACATGAAATCACCAGGTATAAAATTAGACCAAATACCT
85 AATTTTATTACCACTGTAATTAACATGAAATCACCAGGTATAAAATTAGACCAAATACCT
*****

186 CTATTTGTATGAGCGGTAGTAATTACAGCAGTGCTTTTACTATTATCTCTTCCTGTTTTA
72 CTATTTGTATGAGCGGTAGTAATTACAGCAGTGCTTTTACTATTATCTCTTCCTGTTTTA
134 CTATTTGTATGAGCGGTAGTAATTACAGCAGTGCTTTTA-----
85 CTATTTGTATGAGCGGTAGTAATTACAGCAGTGCTTTTACTATTATGGTCTTCCTGTTTT
*****

```

B

Sample #	Park Code	Site	[Hg] (ppb)	% Sequence Similarity
186	ACAD	Eagle Lake	209	100
72	CACO	Gull Pond	70	100
134	CACO	Duck Pond	171	99.5
85	CACO	Great Pond	186	100

Figure 5: A. *Macromia illinoensis* nucleotide sequence alignments. B. Quantification of part A. Park codes are as follows: CACO – Cape Cod National Seashore, ACAD – Acadia National Park.

A

```

46      GGTACTGGTTGAACTGTTTATCCCCCTTTGGCTGGTGCAATTGCCCATGCAGGAGCATCA
53      GGAACAGGTTGAACAGTATACCCTCCTTTGGCTGGTGCAATTGCCCATGCAGGAGCATCA
141     GGTACAGGATGAACAGTTTATCCCCCTTTGGCTGGTGCAATTGCCCATGCAGGAGCATCA
199     -----TTTGGCTGGTGCAATTGCCCATGCAGGAGCATCA
213     -----CCCCCTTTGGCTGGTGCAATTGCCCATGCAGGAGCATCA
118     -----TGGTGCAATTGCCCATGCAGGAGCATCA
          *****

46      GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
53      GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
141     GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
199     GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
213     GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
118     GTAGATCTAACTATTTTTTCTTTACATTTAGCTGGAGTATCTTCAATTCTTGGTGCTATT
          *****

46      AATTTTATTACTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
53      AATTTTATTACTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
141     AATTTTATTACTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
199     AATTTTATTACTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
213     AATTTTATTACTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
118     AATTTTATTAGTACAACAATTAATATAAAGTCACCAGGAATAAAGATAGATCAAATACCT
          *****

46      TTATTTGTATGAGCTGTTGTAATTACAGCTGTACTFTTATTATTATCATTACCTGTTTGG
53      TTATTTGTATGAGCTGTTGTAATTACAGCTGTACTFTTATTATTATCATTACCTGTTTGG
141     TTATTTGTATGAGCTGTTGTAATTACAGCTGTACTFTTATTATTATCATTACCTGTTTGG
199     TTATTTGTATGAGCTGTTGTAATTACAGCTGTACTFTTATTATTATCATTACCTGTTTGG
213     TTATTTGTATGAGCTGTTGTAATTACAGCTGTACTFTTATTATTATCATTACCTGTTTGG
118     TTATTTGTATGAGCTGTTGTAATTACAG-----
          *****

```

B

Sample #	Park Code	Site	[Hg] (ppb)	% Sequence Similarity
46	PIRO	Legion Lake	173	100
53	SACN	Phipps Landing	96	100
141	ACAD	Eagle Lake	209	100
199	ACAD	Lakewood	130	100
213	ACAD	Schoodic	732	100
118	GRPO	Poplar Creek	140	99.5

Figure 6: A. *Aeshna umbrosa* nucleotide sequence alignments. B. Quantification of part A. Park codes are as follows: PIRO – Pictured Rocks National Lakeshore, SACN – Saint Croix National Scenic Riverway, GRPO – Grand Portage National Monument, ACAD – Acadia National Park

A

```

140   AGGAACTGGATGAACTGTTTATCCCCCTTTAGCAGGAGCTATTGCTCATGCTGGTGCATC
82    -GGAACAGGATGAACAGTTTATCCCCCTTTAGCAGGAGCTATTGCTCATGCTGGTGCATC
204   -----TTTAGCAGGAGCTATTGCTCATGCTGGTGCATC
                *****

140   AGTAGACTTAACTATTTTTTCCTTACACCTTGCAGGGGTATCTTCAATTCAGGAGCTAT
82    AGTAGACTTAACTATTTTTTCCTTACACCTTGCAGGGGTATCTTCAATTCAGGAGCTAT
204   AGTAGACTTAACTATTTTTTCCTTACACCTTGCAGGGGTATCTTCAATTCAGGAGCTAT
                *****

140   TAATTTTATTACAACAGTAATTAATATAAAATCACCAGGTATAAAAAATAGATCAAATACC
82    TAATTTTATTACAACAGTAATTAATATAAAATCACCAGGTATAAAAAATAGATCAAATACC
204   TAATTTTATTACAACAGTAATTAATATAAAATCACCAGGTATAAAAAATAGATCAGATACC
                *****

140   ACTATTTGTATGAGCGGTAGTAATTAATGCTGTTCTA-----
82    ACTATTTGTATGAGCGGTAGTAATTAATGCTGTTCTATTACTACTATCATTACCTGTTTT
204   ACTATTTGTATGAGCGGTAGTAATTAATGCTGTT-----
                *****

```

B

Sample #	Park Code	Site	[Hg] (ppb)	% Sequence Similarity
140	ACAD	Jordan Pond	145	100
82	CACO	Gull Pond	70	100
204	ACAD	Eagle Lake	209	99.5

Figure 7: A. *Celithemis elisa* nucleotide sequence alignments. B. Quantification of part A. Park codes are as follows: CACO – Cape Cod National Seashore, ACAD – Acadia National Park

### Amino Acid Sequence Alignments

Amino acid sequence alignments were performed on samples shown by the nucleotide sequence alignments to be mutated (Figure 8). Samples were compared to one non-mutated representative sample of the same species. Two samples were shown to possess nucleotide mutations that translated into amino acid mutations.

A  
65 LAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKMDQMP  
73 LAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKLDQMP  
\*\*\*\*\*:\*\*\*\*\*  
1 53

B  
59 GTGWTVYPPLAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKMDQMP  
40 GTGWTVYPPLAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKMDQMP  
\*\*\*\*\*  
1 60

C  
72 LAGTIAHAGSSVDLTIFSLHLAGVSSILGAINFITTVINMKSPGMKLDQMP  
134 LAGTIAHAGSSVDLTIFSLHLAGVSSILGAINFITTVINMKSPGMKLDQMP  
\*\*\*\*\*  
1 51

D  
118 GAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKMDQMP  
46 GAIAHAGASVDLTIFSLHLAGVSSILGAINFITTTINMKSPGMKMDQMP  
\*\*\*\*\*  
1 49

E  
82 LAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTVINMKSPGMKMDQMP  
204 LAGAIAHAGASVDLTIFSLHLAGVSSILGAINFITTVINMKSPGMKLDQMP  
\*\*\*\*\*:\*\*\*\*  
1 51

F


Sample #	Species	Park Code	Site	[Hg] (ppb)	Amino Acid Changes
65	<i>G. exilis</i>	VOYA	Peary Lake	76	L47M
59	<i>B. janata</i>	VOYA	Peary Lake	76	None
134	<i>M. illinoensis</i>	CACO	Duck Pond	171	None
118	<i>A. umbrosa</i>	GRPO	Poplar Creek	140	None
204	<i>C. elisa</i>	ACAD	Eagle Lake	209	M47L

Figure 8: A – E Amino acid sequence alignments for *G. exilis*, *B. janata*, *M. illinoensis*, *A. umbrosa*, and *C. elisa*. Numbers in red represent amino acid position. F. Quantification of mutated samples. Park codes are as follows: VOYA – Voyageurs National Park, ACAD – Acadia National Park, CACO – Cape Cod National Seashore, GRPO – Grand Portage National Monument

### Protein Structure Models

Models of protein structure for the samples with mutated amino acid sequences were compared to models generated from the non-mutated amino acid sequences of samples of the same species (Figure 9). Both non-mutated sequences generated a structure consisting of a single alpha helix, while both mutated sequences generated a structure consisting of a double alpha helix.

A



B




Figure 9: Protein Structure Models. Samples are rainbow colored starting at the N-terminus, in red. A. *G. exilis* samples 73 (non-mutated, left) and 65 (mutated, right). B. *C. elisa* samples 82 (non-mutated, left) and 204 (mutated, right).

## DISCUSSION AND CONCLUSIONS

The cytochrome c oxidase I (COI) gene is widely used for DNA barcoding and Figure 2 illustrates why this gene is so effective for species identification. In parts A and B, the nucleotide and amino acid sequences of five different dragonfly species are aligned. Part A shows that while the genes maintain enough sequence identity to be aligned, there are many variations in the nucleotide sequence. Despite these variations, the redundancy of the genetic code ensures nearly identical amino acid sequences, shown in part B. This is expected because the COI protein structure would need to be conserved for it to function properly during oxidative phosphorylation.

Despite COI being well suited for DNA barcoding, only 41% of samples were successfully identified to the species level due to poor sequencing. It is suspected that the PCR primers may not have been optimal for all genus or species. Primers used were developed relatively recently as an alternative for the traditional primers to yield a 313 bp PCR fragment rather than a 658 bp PCR fragment to enable easier sequencing. The 313 bp fragment is the last 313 bp of the 658 bp fragment.<sup>34</sup> It is advisable in the future to conduct a comparative study of the efficiency of the two primer sets as the original comparison was evaluated in fish, so perhaps the primers are not well suited for dragonflies.

Many studies have documented that mercury is able to bioaccumulate in dragonflies, however the effects of mercury on dragonflies has not been well investigated.<sup>39, 40, 41</sup> Mercury has been shown in other organisms to act as a mutagen. In a study using mouse embryonic fibroblasts transfected with the *lacZ* reporter gene, mercury

exposure was shown to cause DNA mutations at a concentration of 50 ppb. Mutation frequencies at 100 ppb and 150 ppb were not significantly different from each other, but were higher than at 50 ppb. Mercury concentrations higher than 150 ppb were not tested<sup>26</sup>. The potential of mercury to cause DNA damage at these concentrations in other organisms is significant because the State of Maine fish consumption advisory concentration of mercury is 200 ppb.<sup>42</sup> While there are no published thresholds to indicate concentrations of mercury that would have species, population, or higher trophic level consumption implications for dragonflies, because they are known vectors for mercury biomagnification the effects of mercury on these organisms is worth investigating.

To attempt to find any mutations in the COI gene corresponding to mercury exposure, the nucleotide sequences were aligned from dragonfly species found to be living in habitats associated with relatively high and relatively low dragonfly larvae mercury concentrations (at least one habitat <100 ppb dw and one >200 ppb dw). Five species fit this description: *Gomphus exilis*, *Basiaeschna janata*, *Macromia illinoiensis*, *Aeshna umbrosa*, and *Celithemis elisa*. It was hypothesized that samples found at sites that would subject them to a relatively higher concentration of mercury would have more mutations than samples found at sites that would subject them to a relatively lower concentration of mercury. As shown in Figures 3 – 7, each species had one representative member with at least one mutation. All mutations detected were point mutations. This is consistent with the findings of a human population study that studied the DNA repair efficiency of lymphocytes taken from factory workers occupationally exposed to

mercury. The study concluded that mercury exposure impairs base excision repair mechanisms.<sup>22</sup>

With the exception of *C. elisa* sample 204, the rest of the mutated samples were from sites that contained dragonflies with less than 200 ppb of mercury, on average. These findings agree with the previously mentioned mouse model study in the sense that DNA mutations did occur at in dragonflies at the same mercury concentrations that they occurred in mice, but mutation frequency did not show a trend of increasing as mercury concentration increased, which did not support the hypothesis. The chi-squared statistic for this data was calculated to be 2.263 with a *p*-value of 0.323, which is not significant at  $p < 0.05$ . While the result of most sequence variation occurring at mercury concentrations less than 200 ppb was unexpected, it suggests that genetic variation within a species is reduced in areas with a higher mercury concentration. As there was only one sample site tested with a mercury concentration significantly above 200 ppb, this cannot be definitively concluded.

Point mutations in the coding region of a nucleotide sequences are the most dangerous if they lead to changes in the amino acid sequence, which could change in the protein structure to reduce or inhibit function. To test for this, the amino acid sequences of the samples with mutated nucleotide sequences were aligned with the amino acid sequences of samples of the same species whose nucleotide sequences weren't mutated. For three samples, the mutation(s) present in the nucleotide sequence caused no change in the amino acid sequence (Figure 8), but for two samples these mutations led to mutations in the amino acid sequence. *G. exilis* sample 65 underwent a Leucine to Methionine mutation, and *C. elisa* sample 204 underwent a Methionine to Leucine mutation. Both

mutations occurred at the same locus. Looking back at the nucleotide sequences for these two samples, they both underwent mutations that led to the incorporation of a guanine. It is possible this guanine is 8-oxo-dGTP, in which case this mutation would be the result of oxidative damage.<sup>14, 15, 16</sup>

To determine if these amino acid mutations altered the protein structure of COI, models of the structures were generated, shown in Figure 9. Both of the non-mutated control samples had a secondary structure consisting of a single alpha-helix. Both mutated samples had altered structures consisting of a large alpha-helix accompanied by a smaller alpha-helix. It is interesting that despite these mutations occurring at the same locus and being essentially the reverse of each other, that they both caused the same change in secondary structure. It is possible that this change in structure could impact COI function.

These results show that while more mutations were not found in samples from sites with relatively higher concentrations of mercury, some mutations that did occur caused alterations to the amino acid sequence and protein structure of COI. Despite mercury being a known mutagen, because these dragonflies were collected from the environment we cannot be certain that mercury exposure was the cause of the detected mutations. They could have been exposed to another, un-tested mutagen before being collected. It is also possible that these mutations are just contributing to the natural variation present between individuals and simply happened over time.

Future experiments need to be performed to draw more definitive conclusions from these results. Analysis should be performed on COI mutants with altered protein structure to determine if they are still functional and in what capacity. To determine if

mercury is actually reducing genetic variation at concentrations over 200 ppb, a greater range of concentrations need to be analyzed than those examined in this study. If sampling sites at this range are not available, this could be done using artificial exposure of dragonflies to mercury in a lab. Artificial exposure would also help to determine if the mutations shown are actually being caused by mercury because researchers could limit exposure to other pollutants.

As this was the first study of its kind (to our knowledge), dragonfly leg tissue was used for mutation analysis so taxonomic identification of samples would still be possible in the event DNA barcoding failed. In the future results may be improved by utilizing abdomen tissue as it has been shown to be the body region where mercury accumulates the most in dragonflies.<sup>39</sup> However, it should be noted that because mitochondria are maternally inherited, and COI is a mitochondrial gene, detected mutation may have been maternally inherited, and in this case would not be specific to a particular body region.

This study was conducted with a very small sample size, which also impacts the ability to draw strong conclusions. The samples are randomly collected from the field and species is identified later, so ensuring the collection of a certain number of species is impossible. However only one year was examined for this study. Examining multiple years' worth of data could help to increase the sample size and thus strengthen the conclusions.

## BIBLIOGRAPHY

1. Gworek, B., Bemowska-Kalabun, O., Kijewska, M., & Wrzosek-Jakubowska, J. (2016). Mercury in marine and oceanic waters—a review. *Water, Air, & Soil Pollution*, 227 (10), 1-19. DOI:10.1007/s11270-016-3060-3
2. Jozef M Pacyna, Oleg Travnikov, Francesco De Simone, Ian M Hedgecock, Kyrre Sundseth, Elisabeth G Pacyna, . . . Karin Kindbom. (2016). Current and future levels of mercury atmospheric pollution on a global scale. *Atmospheric Chemistry and Physics*, 16(19), 12495-12511. DOI:10.5194/acp-16-12495-2016
3. Nriagu, J., & Becker, C. (2003). Volcanic emissions of mercury to the atmosphere: Global and regional inventories. *Science of the Total Environment*, 304(1), 3-12. DOI:10.1016/S0048-9697(02)00552-1
4. Sundseth, K., Pacyna, J., Pacyna, E., Pirrone, N., & Thorne, R. (2017). Global sources and pathways of mercury in the context of human health. *International Journal of Environmental Research and Public Health*, 14(1), 105. DOI:10.3390/ijerph14010105
5. Pirrone, N., & Mason, R. (2009). *Mercury Fate and Transport in the Global Atmosphere*. Dordrecht, The Netherlands: Springer.
6. Kabata-Pendias, A., & Mukherjee, A. B. (2007). *Trace elements from soil to human*. Springer Science & Business Media.
7. Crespo-López, M. E., Macêdo, G. L., Pereira, S. I. D., Arrifano, G. P. F., Picanço-Diniz, D. L. W., Nascimento, José Luiz M do, & Herculano, A. M. (2009). Mercury and human genotoxicity: Critical considerations and possible molecular mechanisms. *Pharmacological Research*, 60(4), 212-220. DOI:10.1016/j.phrs.2009.02.011
8. United States Environmental Protection Agency. (1997) Mercury study report to congress: health effects of mercury and mercury compounds. EPA-4562/R-97-007.
9. National research Council. (2000). *Toxicological effects of methylmercury*. National Academies Press.
10. C. Baird, & M. Cann (2004) *Environmental Chemistry* W.H. Freeman, New York
11. Leonard, S. S., Harris, G. K., & Shi, X. (2004). Metal-induced oxidative stress and signal transduction. *Free Radical Biology and Medicine*, 37(12), 1921-1942. DOI:10.1016/j.freeradbiomed.2004.09.010
12. Halliwell, B., & Gutteridge, J. (1984). Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochemical journal*, 219(1), 1.

13. Chen, F., & Shi, X. (2002). Intracellular signal transduction of cells in response to carcinogenic metals. *Critical reviews in oncology/hematology*, 42(1), 105-121. DOI:10.1016/S1040-8428(01)00211-6
14. Fasullo, M., & Endres, L. (2015). Nucleotide salvage deficiencies, DNA damage and neurodegeneration. *International Journal of Molecular Sciences*, 16(5), 9431-9449. DOI:10.3390/ijms16059431
15. Furda, A. M., Marrangoni, A. M., Lokshin, A., & Van Houten, B. (2012). Oxidants and not alkylating agents induce rapid mtDNA loss and mitochondrial dysfunction. *DNA repair*, 11(8), 684-692. DOI:10.1016/j.dnarep.2012.06.002
16. Scheibye-Knudsen, M., Fang, E. F., Croteau, D. L., Wilson, D. M., & Bohr, V. A. (2015). Protecting the mitochondrial powerhouse. *Trends in cell biology*, 25(3), 158-170. DOI:10.1016/j.tcb.2014.11.002
17. Pursell, Z. F., McDonald, J. T., Mathews, C. K., & Kunkel, T. A. (2008). Trace amounts of 8-oxo-dGTP in mitochondrial dNTP pools reduce DNA polymerase  $\gamma$  replication fidelity. *Nucleic acids research*, 36(7), 2174-2181. DOI:10.1093/nar/gkn062
18. Sener, G., Sehirli, A., Dulger, G. (2003). Melatonin protects against mercury(II)-induced oxidative tissue damage in rats. *Pharmacology & Toxicology*, 93(6), 290-296. DOI:10.1111/j.1600-0773.2003.pto930607.x
19. Lund, B. O., Miller, D. M., & Woods, J. S. (1993). Studies on Hg (II)-induced H<sub>2</sub>O<sub>2</sub> formation and oxidative stress in vivo and in vitro in rat kidney mitochondria. *Biochemical pharmacology*, 45(10), 2017-2024. DOI:10.1016/0006-2952(93)90012-L
20. Mahboob, M., Shireen, K. F., Atkinson, A., & Khan, A. T. (2001). Lipid peroxidation and antioxidant enzyme activity in different organs of mice exposed to low level of mercury. *Journal of Environmental Science and Health, Part B*, 36(5), 687-697. DOI:10.1081/PFC-100106195
21. Nava, M., Romero, F., Quiroz, Y., Parra, G., Bonet, L., & Rodríguez-Iturbe, B. (2000). Melatonin attenuates acute renal failure and oxidative stress induced by mercuric chloride in rats. *American Journal of Physiology-Renal Physiology*, 279(5), 910-918.
22. Cebulska-Wasilewska, A., Panek, A., Żabiński, Z., Moszczyński, P., & Au, W. W. (2005). Occupational exposure to mercury vapour on genotoxicity and DNA repair. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 586(2), 102-114. DOI:10.1016/j.mrgentox.2005.06.009
23. Hartwig, A., Asmuss, M., Blessing, H., Hoffmann, S., Jahnke, G., Khandelwal, S., ... & Bürkle, A. (2002). Interference by toxic metal ions with zinc-dependent proteins involved in maintaining genomic stability. *Food and Chemical Toxicology*, 40(8), 1179-

1184. DOI:10.1016/S0278-6915(02)00043-1

24. Sekowski, J. W., Malkas, L. H., Wei, Y., & Hickey, R. J. (1997). Mercuric ion inhibits the activity and fidelity of the human cell DNA synthesome. *Toxicology and applied pharmacology*, 145(2), 268-276. DOI:10.1006/taap.1997.8185

25. Li, Y., Jiang, Y., & Yan, X. P. (2006). Probing Mercury Species– DNA Interactions by Capillary Electrophoresis with On-Line Electrothermal Atomic Absorption Spectrometric Detection. *Analytical chemistry*, 78(17), 6115-6120. DOI:10.1021/ac060644a

26. Schurz, F., Sabater-Vilar, M., & Fink-Gremmels, J. (2000). Mutagenicity of mercury chloride and mechanisms of cellular defence : The role of metal-binding proteins. *Mutagenesis*, 15(6), 525-530.

27. Munday, R. (1994) Bioactivation of thiols by one-electron oxidation. *Advances in Pharmacology*, 27, 237-270.

28. Lin, T., & Yang, M. S. (2007). Benzo[a]pyrene-induced elevation of GSH level protects against oxidative stress and enhances xenobiotic detoxification in human HepG2 cells. *Toxicology*, 235(1–2), 1-10. DOI: 10.1016/j.tox.2007.03.002

29. Dringen, R. (2000). Glutathione metabolism and oxidative stress in neurodegeneration. *European Journal of Biochemistry*, 267(16), 4903-4903. DOI:10.1046/j.1432-1327.2000.01651.x

30. Deneke, S. M., & Fanburg, B. L. (1989). Regulation of cellular glutathione. *American Journal of Physiology-Lung Cellular and Molecular Physiology*, 257(4), 163-173.

31. Richman, P. G., & Meister, A. (1975). Regulation of gamma-glutamyl-cysteine synthetase by nonallosteric feedback inhibition by glutathione. *Journal of Biological Chemistry*, 250(4), 1422-1426.

32. Kachur, A. V., Koch, C. J., & Biaglow, J. E. (1998) Mechanism of copper-catalyzed oxidation of glutathione. *Free Radic. Res.* 28 (3), 259-269. DOI:10.3109/10715769809069278

33. Paul D. N. Hebert, Alina Cywinska, Shelley L. Ball, & Jeremy R. deWaard. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1512), 313-321. DOI:10.1098/rspb.2002.2218

34. Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., . . . Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: Application for

- characterizing coral reef fish gut contents. *Frontiers in Zoology*, 10(1), 34.  
DOI:10.1186/1742-9994-10-34
35. Cox, A. J., & Hebert, P. D. (2001). Colonization, extinction, and phylogeographic patterning in a freshwater crustacean. *Molecular Ecology*, 10(2), 371-386.  
DOI:10.1046/j.1365-294X.2001.01188.x
36. Wares, J. P., & Cunningham, C. W. (2001). Phylogeography and historical ecology of the North Atlantic intertidal. *Evolution*, 55(12), 2455-2469.
37. Tiziana Venesio, Antonella Balsamo, Edoardo Errichiello, Guglielmina N Ranzani, & Mauro Risio. (2013). Oxidative DNA damage drives carcinogenesis in MUTYH-associated-polyposis by specific mutations of mitochondrial and MAPK genes. *Modern Pathology*. 26(10), 1371-1381. DOI:10.1038/modpathol.2013.66
38. Eagles-Smith, C.A., Nelson, S.J., Willacker, J.J., Jr., Flanagan Pritz, C.M., and Krabbenhoft, D.P. (2016). Dragonfly Mercury Project—A citizen science driven approach to linking surface-water chemistry and landscape characteristics to biosentinels on a national scale: U.S. Geological Survey Fact Sheet 2016-3005, 4 p.  
DOI:10.3133/fs20163005
39. Buckland-Nicks, A., Hillier, K. N., Avery, T. S., & O'Driscoll, N. J. (2014). Mercury bioaccumulation in dragonflies (odonata: Anisoptera): Examination of life stages and body regions. *Environmental Toxicology and Chemistry*, 33(9), 2047-2054.  
doi:10.1002/etc.2653
40. Haro, R. J., Bailey, S. W., Northwick, R. M., Rolfhus, K. R., Sandheinrich, M. B., & Wiener, J. G. (2013). Burrowing dragonfly larvae as biosentinels of methylmercury in freshwater food webs. *Environmental Science & Technology*, 47(15), 8148.
41. Zhang, Z., Song, X., Wang, Q., & Lu, X. (2012). Mercury bioaccumulation and prediction in terrestrial insects from soil in huludao city, northeast china. *Bulletin of Environmental Contamination and Toxicology*, 89(1), 107-112. doi:10.1007/s00128-012-0649-0
42. Maine Bureau of Health, Fish Tissue Action Levels. Feb. 20, 2001.

## APPENDIX

Standard Operating Procedure for DNA Barcoding of Larval Dragonflies

Megan Little

University of Maine

Karen James, Michael Marion - MDI Biological Laboratory

Hamish Grieg, Sarah J. Nelson – University of Maine

A contribution to the Dragonfly Mercury Project

## A. Preparing the Tissue Samples

### Materials and Equipment:

- Preserved<sup>1</sup> dragonflies
- Glass petri dish
- Kimwipes<sup>®</sup>
- Forceps
- Scissors
- 1.5 mL microcentrifuge tubes
- Flame source (Bunsen burner, lighter, etc.)

### Procedure:

1. Sterilize all instruments and the glass petri dish by passing them over a flame before beginning the dissections.
2. Working with one dragonfly at a time, cut all the legs off one side of the dragonfly using the scissors.
3. Place all legs into one microcentrifuge tube labeled with the sample ID and date. If not moving directly to part C, freeze sample in -20°C freezer.
4. Sterilize all instruments used in the dissection, including the glass dish. Replace the Kimwipe<sup>®</sup> on the work surface.
5. Repeat procedure for all dragonflies being analyzed.

## B. DNA Extraction

### Materials and Equipment:

- Qiagen DNeasy Blood and Tissue Kit
- 1.5 mL microcentrifuge tubes
- Heat block or oven
- Microcentrifuge
- Vortex
- Pipettes and sterile, disposable pipette tips (ideally filter tips)
- Disposable plastic pestles

### Procedure:

1. Add 180  $\mu$ L of Buffer ATL and 20  $\mu$ L Proteinase K to each sample and vortex for 5 seconds. Incubate overnight at 55°C.
2. Vortex each sample for 15 seconds, add 200  $\mu$ L Buffer AL, then grind tissue with a new plastic pestle. Vortex again and incubate at 55°C for 20 minutes.
3. Add 200  $\mu$ L of ethanol to each sample and vortex.
4. Working with each sample separately, pipette the entire mixture volume into a DNeasy Mini Spin Column placed in a 2 mL collection tube. Centrifuge at 8000 rpm for 1 minute in a microcentrifuge.
5. Discard the flow-through and replace the collection tube. Pipette 500  $\mu$ L Buffer AW1 into the column and centrifuge at 8000 rpm for 1 minute.

---

<sup>1</sup> Dragonflies may be preserved by either freezing or submersion in 95% ethanol.

6. Discard the flow-through and replace the collection tube. Pipette 500  $\mu\text{L}$  Buffer AW2 into the column and centrifuge at 14,000 rpm for 3 minutes.
7. Discard the flow-through and discard the collection tube, placing the column into a 1.5 mL microcentrifuge tube.
8. Pipette 100  $\mu\text{L}$  Buffer AE into the column and incubate at room temperature for 1 minute. Centrifuge at 8000 rpm for 1 minute to elute the DNA.
9. Pipette 50  $\mu\text{L}$  of Buffer AE into the column and incubate at 55°C for 5 minutes. Centrifuge at 8000 rpm for 1 minute to elute the remaining DNA.
10. Discard the column and quantify the DNA using the nanodrop.
11. Store the purified DNA in a -20°C freezer if not moving directly to part D.

### C. PCR

#### Materials and Equipment:

- m1COIintF and jgHCO2198 primers
- Sigma RedTaq ReadyMix
- Purified DNA
- Microcentrifuge or mini-microcentrifuge
- Pipettes and sterile, disposable pipette tips (ideally filter tips)
- Thermocycler

#### Procedure:

1. Use <http://www.idtdna.com/Calc/resuspension> to calculate how much nuclease-free water is needed to resuspend your new primers to form a 100  $\mu\text{M}$  stock solution. Make an aliquot of 10  $\mu\text{M}$  working solution.
2. Make your master-mix in a 1.5 mL microcentrifuge tube using the following component volumes per reaction, including a negative control and 2 extra reactions: 12.5  $\mu\text{L}$  Sigma RedTaq ReadyMix, 10.5  $\mu\text{L}$  nuclease-free water, 0.5  $\mu\text{L}$  forward primer, and 0.5  $\mu\text{L}$  reverse primer. Vortex then spin mixture in mini-microcentrifuge, then keep on ice.
3. Add 24  $\mu\text{L}$  of master-mix to 1 0.2 mL PCR tube per reaction.
4. Add 1  $\mu\text{L}$  of DNA to each tube. Use 1  $\mu\text{L}$  nuclease-free water for the negative control.
5. Spin tubes in the mini-microcentrifuge then place them in the thermal cycler. Run samples on an amplification program which should have the following conditions:
  - a. 16 cycles of the following:
    - i. 95°C for 10 seconds
    - ii. 62°C for 30 seconds, -1° per cycle
    - iii. 72°C for 60 seconds
  - b. 25 cycles of the following:
    - i. 95°C for 10 seconds
    - ii. 46°C for 30 seconds
    - iii. 72°C for 60 seconds
  - c. Final hold at 4°C
6. Samples can be frozen in a -20°C freezer if not moving directly on to part E.

## D. Gel Electrophoresis

### Materials and Equipment:

- PCR products
- Invitrogen Low DNA Mass Ladder
- Pipettes and sterile, disposable pipette tips (ideally filter tips)

### Large 1.5% Agarose Gel (Maximum 40 wells):

- 150 mL TBE Buffer
- 2.25 g Agarose
- 2.5  $\mu$ L Ethidium Bromide

### Small 1.5% Agarose Gel (Maximum 24 wells):

- 60 mL TBE Buffer
- 0.9 g Agarose
- 2.0  $\mu$ L Ethidium Bromide

### Procedure:

1. Combine TBE buffer and agarose in a flask.
2. Heat in a microwave in 30 or 60 second intervals depending on if making a small or large gel. Heat until mixture boils and appears clear.
3. Cool flask until it can be touched for three seconds on any side.
4. Add ethidium bromide and gently swirl mixture. Pour into gel rig. Let sit at least half an hour.
5. Pipette 5  $\mu$ L of Invitrogen Low DNA Mass Ladder into the first well. Pipette 15  $\mu$ L of each PCR product into its own well, making sure to include the negative control.
6. Run the gel for 30 minutes at voltage specified above then view using a UV transilluminator with a mounted camera. If you do not see bands for your PCR products, go back and troubleshoot your PCR. If your PCR products show bands, you can proceed.
7. Samples can be frozen in a  $-20^{\circ}\text{C}$  freezer if not moving directly on to part F.

## E. Sample Clean-Up and Sequencing at the University of Maine's DNA Sequencing Facility

### Materials and Equipment:

- Affymetrix ExoSAP-IT
- 0.2 mL microcentrifuge tubes
- 10  $\mu$ M PCR primers
- Pipettes and sterile, disposable pipette tips (ideally filter tips)
- UV Transilluminator

### Procedure:

1. Add 2  $\mu$ L of Affymetrix ExoSAP-IT to one 0.2 mL microcentrifuge tube for each PCR product. Keep tubes on ice.
2. Add 5  $\mu$ L of a PCR product to each tube and spin briefly in a mini-microcentrifuge.

3. Place tubes in a thermal cycler and run on the ExoSAP program. The product can either be frozen in a -20°C freezer or you can proceed.
4. In a new 0.2 mL microcentrifuge tube, dilute 1 µL of each PCR product with enough nuclease-free water to yield a final DNA concentration of 10 ng/µL.
5. Briefly spin tubes in the mini-microcentrifuge and deliver to the sequencing lab.

#### F. Sample Identification

##### Procedure:

1. Once sequence files returned, typically in “.abi” format, open them in a DNA editing software (e.g. 4Peaks for Mac OSX) then:
  - a. Visually scan sequences for sequence quality.
  - b. Trim the ends to remove poor sequence quality.
  - c. For sequences from the reverse primer, “Flip Sequence” (reverse complement).
  - d. Highlight and copy (ctrl-c) sequence.
2. Go to the Barcode of Life Database (<http://boldsystems.org>) and click on the “Identification” tab.
3. While under the “Animal Identification [COI]” tab, paste (ctrl-v) your sequence into the text box.
4. If no match is found, repeat selecting less stringent search options.
5. Check that the search results for the forward and reverse sequences from each specimen match.

*Note: All standard lab safety procedures should be followed throughout, including, but not limited to, use of safety glasses or goggles, gloves, and other personal protective equipment; the separation of the sterilizing flame from areas in which ethanol is being used; and storage of reagents and chemicals in an appropriate manner.*

## AUTHOR'S BIOGRAPHY

Megan Little was born in Bangor, Maine on May 31, 1995. She was raised in Calais, Maine and graduated valedictorian of her class from Calais High School in 2013. Megan is a double major in molecular and cellular biology and biochemistry. In the fall she will continue her education at Marshall University working towards obtaining a Master's degree in forensic science with a concentration in DNA analysis.